

MIXED FRACTIONAL DIFFERENTIATION OPERATORS IN HÖLDER SPACES
DEFINED BY USUAL HÖLDER CONDITION

Mamatov Tulkin, Sabirova Rano, Barakaev Dilshod

**Teachers of Department of Higher mathematics, Bukhara Technological
Institute of Engineering
706030, Uzbekistan, Bukhara Region, Bukhara Street, K. Murtazayev, 15**

ABSTRACT

We study mixed fractional derivative in Marchaud form of function of two variables in Hölder spaces of different orders in each variables. The main interest being in the evaluation of the latter for the mixed fractional derivative in the cases Hölder class defined by usual Hölder condition.

АННОТАЦИЯ

Мы изучаем смешанную дробную производную в форме Маршо от двух переменных в пространствах Гельдера.

Keywords: functions of two variables, fractional derivative of Marchaud form, mixed fractional derivative, mixed fractional integral, Hölder space.

Ключевые слова: функции двух переменных, дробный дифференциал форме Маршо, смешанный дробный производный, смешанный дробный интеграл, пространства Гельдера.

Introduction

The classical result of G.Hardi and D.Littlewood (1928, see [1, §3]) is known that the fractional integral $(I_{a+}^{\alpha} f)(x) = \Gamma^{-1}(\alpha) (t_+^{\alpha-1} * f)(x)$, $0 < \alpha < 1$ maps isomorphically the space $H_0^{\lambda}([0, 1])$ of Hölder order $\lambda \in (0, 1)$ functions with a condition $f(0) = 0$ on a similar space of a higher order $\lambda + \alpha$ provided that

$\lambda + \alpha < 1$. Further, this result was generalized in various directions: a space with a power weight, generalized Hölder spaces, spaces of the Nikolsky type, etc. A detailed review of these and some other similar results can be found in [1].

In the multidimensional case, the statement about the properties of a map in Hölder spaces for a mixed fractional Riemann – Liouville integral was studied in [2] - [7].

Mixed fractional derivatives form Marchaud

$$\begin{aligned} (D_{a^+, c^+}^{\alpha, \beta} \phi)(x, y) = & \frac{\phi(x, y)}{\Gamma(1-\alpha)\Gamma(1-\beta)(x-a)^\alpha(y-c)^\beta} + \\ & \int_y \frac{\phi(x, y) - \phi(t, \tau)}{(x-t)^{1+\alpha}(y-\tau)^{1+\beta}} dt d\tau, \quad x > a, y > c, \end{aligned} \quad (1)$$

were not studied in the Hölder space. This paper is devoted to the study of the properties of a map in Hölder spaces, defined by the usual Hölder condition for functions of two variables.

Consider the operator (1) in a rectangle $Q = \{(x, y): 0 < x < b, 0 < y < d\}$.

For a continuous function $\phi(x, y)$ on R^2 we introduce the notation

$$\begin{aligned} \left(\Delta_h^{1,0} \phi \right)(x, y) &= \phi(x+h, y) - \phi(x, y), \quad \left(\Delta_\eta^{0,1} \phi \right)(x, y) = \phi(x, y+\eta) - \phi(x, y), \\ \left(\Delta_{h,\eta}^{1,1} \phi \right)(x, y) &= \phi(x+h, y+\eta) - \phi(x+h, y) - \phi(x, y+\eta) + \phi(x, y), \end{aligned}$$

so that

$$\phi(x+h, y+\eta) = \left(\Delta_{h,\eta}^{1,1} \phi \right)(x, y) + \left(\Delta_h^{1,0} \phi \right)(x, y) + \left(\Delta_\eta^{0,1} \phi \right)(x, y) + \phi(x, y). \quad (2)$$

Everywhere in the sequel by C, C_1, C_2 etc we denote positive constants which may have different values in different occurrences and even in the same line.

Definition 1. Let $\lambda, \gamma \in (0, 1]$. We say that $\phi \in H^{\lambda, \gamma}(Q)$, if

$$|\phi(x_1, y_1) - \phi(x_2, y_2)| \leq C_1 |x_1 - x_2|^\lambda + C_2 |y_1 - y_2|^\gamma \quad (3)$$

for all $(x_1, y_1), (x_2, y_2) \in Q$. Condition (3) is equivalent to the couple of the separate conditions

$$\left| \binom{1,0}{\Delta_h \phi}(x,y) \right| \leq C_1 |h|^\lambda, \quad \left| \binom{0,1}{\Delta_\eta \phi}(x,y) \right| \leq C_2 |\eta|^\nu \quad (4)$$

uniform with respect to another variable.

Note that

$$\phi(x,y) \in H^{\lambda,\nu} \Rightarrow \left| \binom{1,1}{\Delta_{h,\eta} \phi}(x,y) \right| \leq C_\theta |h|^{\theta\lambda} |\eta|^{(1-\theta)\nu} \leq C \min\{|h|^\lambda, |\eta|^\nu\}, \quad \text{where } \theta \in [0,1]. \quad (5)$$

By $H_0^{\lambda,\nu}(Q)$ we define a subspace of functions $f \in H_0^{\lambda,\nu}(Q)$, vanishing at the boundaries $x=0$ and $y=0$ of Q .

A one –dimensional statements

The following statements are known [1]. We use the schemes of the proofs to make the presentation easier for two-dimensional case.

Lemma 1. If $f(x) \in H^{\lambda+\alpha}([0, b])$ and $0 < \lambda, 0 < \alpha + \lambda < 1$, then

$$g(x) = \frac{f(x) - f(0)}{|x|^\alpha} \in H^\lambda([0, b]) \quad \text{and} \quad \|g\|_{H^\lambda} \leq C \|f\|_{H^{\lambda+\alpha}},$$

where C doesn't depend from $f(x)$.

Proof. Let $h > 0; x, x+h \in [0, b]$. We consider the difference

$$|g(x+h) - g(x)| \leq \frac{|f(x+h) - f(x)|}{(x+h)^\alpha} + |f(x) - f(0)| \frac{(x+h)^\alpha - x^\alpha}{x^\alpha (x+h)^\alpha}.$$

Since $f \in H^{\lambda+\alpha}$, we have

$$|f(x+h) - f(x)| \leq C_1 h^{\lambda+\alpha}, \quad |f(x) - f(0)| \leq C_2 x^{\lambda+\alpha}.$$

Using these inequalities we obtain

$$|g(x+h) - g(x)| \leq C_1 \frac{h^{\lambda+\alpha}}{(x+h)^\alpha} + C_2 x^\lambda \frac{(x+h)^\alpha - x^\alpha}{(x+h)^\alpha} = G_1 + G_2.$$

For G_1 we have

$$G_1 = C_1 h^\lambda \left(\frac{h}{x+h} \right)^\alpha \leq C h^\lambda.$$

Let's estimate G_2 , here we shall consider two cases: $x \leq h$ and $x > h$. In the first case, we use inequality $|\sigma_1^\mu - \sigma_2^\mu| \leq |\sigma_1 - \sigma_2|^\mu$, ($\sigma_1 \neq \sigma_2$) and obtain

$$G_2 \leq C_2 \frac{x^\lambda h^\alpha}{(x+h)^\alpha} \leq Ch^\lambda$$

In second case, using $(1+t)^\alpha - 1 \leq \alpha t$, $t > 0$ we have

$$G_2 = C_2 \frac{x^{\lambda+\alpha}}{(x+h)^\alpha} \left| \left(1 + \frac{h}{x}\right)^\alpha - 1 \right| \leq Chx^{\lambda-1} \leq Ch^\lambda$$

which completes the proof.

The Marchaud fractional differentiation operator has a form:

$$(D_{0+}^\alpha f)(x) = \frac{f(x)}{x^\alpha \Gamma(1-\alpha)} + \frac{\alpha}{\Gamma(1-\alpha)} \int_0^x \frac{f(x) - f(t)}{(x-t)^{1+\alpha}} dt, \text{ where } 0 < \alpha < 1. \quad (6)$$

Theorem 1. If $f(x) \in H^{\lambda+\alpha}([0, b])$, $0 < \alpha + \lambda < 1$, that

$$(D_{0+}^\alpha f)(x) = \frac{f(0)}{x^\alpha \Gamma(1-\alpha)} + \psi(x), \quad (7)$$

where $\psi(x) \in H^\lambda([0, b])$ and $\psi(0) = 0$, thus $\|\psi\|_{H^\lambda} \leq C \|f\|_{H^{\lambda+\alpha}}$.

Proof. We present (6) as

$$(D_{0+}^\alpha f)(x) = \frac{f(0)}{x^\alpha \Gamma(1-\alpha)} + \frac{f(x) - f(0)}{x^\alpha \Gamma(1-\alpha)} + \frac{\alpha}{\Gamma(1-\alpha)} \int_0^x \frac{f(x) - f(t)}{(x-t)^{1+\alpha}} dt,$$

receive equality (7), where

$$\psi(x) = \psi_1(x) + \psi_2(x) = \frac{f(x) - f(0)}{x^\alpha \Gamma(1-\alpha)} + \frac{\alpha}{\Gamma(1-\alpha)} \int_0^x \frac{f(x) - f(t)}{(x-t)^{1+\alpha}} dt.$$

Here $\psi_1(x) \in H^\lambda([0, b])$ by Lemme 1. It is enough to show $\psi_2(x) \in H^\lambda([0, b])$.

Let $h > 0$; $x, x+h \in [0, b]$. Let's consider the difference

$$\begin{aligned} \psi_2(x+h) - \psi_2(x) &= \int_0^x \frac{f(x+h) - f(x)}{(x+h-t)^{1+\alpha}} dt + \int_x^{x+h} \frac{f(x+h) - f(t)}{(x+h-t)^{1+\alpha}} dt + \\ &+ \int_0^x [f(x) - f(t)] \left[\frac{1}{(x+h-t)^{1+\alpha}} - \frac{1}{(x-t)^{1+\alpha}} \right] dt = I_1 + I_2 + I_3. \end{aligned}$$

Since $f \in H^{\lambda+\alpha}([0, b])$, then we have for I_1

$$|I_1| \leq Ch^{\lambda+\alpha} \int_0^x (t+h)^{-1-\alpha} dt \leq C_1 h^\lambda .$$

Let's estimate I_2 . We have

$$|I_2| \leq C \int_x^{x+h} (x+h-t)^{\lambda-1} dt = C_2 h^\lambda$$

For I_3 ,

$$|I_3| \leq Ch^\lambda \int_0^{\frac{x}{h}} t^\lambda \left| \frac{1}{(1+t)^{1+\alpha}} - \frac{1}{t^{1+\alpha}} \right| dt \leq C_3 h^\lambda .$$

Finally, it remains to note that $\psi_2(0)=0$, since

$$|\psi_2(x)| \leq C \int_0^x t^{\lambda-1} dt .$$

Main result.

Lemma 2. Let $f(x, y) \in H^{\lambda, \nu}(Q)$, $\alpha < \lambda \leq 1$, $\beta < \nu \leq 1$. Then for the mixed fractional differential operator (1) the representation

$$(D_{0+,0+}^{\alpha,\beta} f)(x, y) = \frac{f(0,0) x^{-\alpha} y^{-\beta} + y^{-\beta} \psi_1(x) + x^{-\alpha} \psi_2(y) + \psi(x, y)}{\Gamma(1-\alpha) \Gamma(1-\beta)}, \quad (8)$$

and

$$|\psi_1(x)| \leq C_1 x^{\lambda-\alpha}, \quad |\psi_2(y)| \leq C_2 y^{\nu-\beta},$$

(9)

$$|\psi(x, y)| \leq C x^{\theta\lambda-\alpha} y^{(1-\theta)\nu-\beta} \quad (10)$$

where

$$\begin{aligned} \psi_1(x) &= \int_0^x [f(x, 0) - f(t, 0)] (x-t)^{-\alpha-1} dt, \\ \psi_2(y) &= y^{-\beta} [f(0, y) - f(0, 0)] + \beta \int_0^y [f(0, y) - f(0, \tau)] (y-\tau)^{-1-\beta} d\tau, \end{aligned}$$

$$i_y \left(\Delta_{x-t, y-\tau}^{1,1} f \right) (t, \tau) \frac{dtd\tau}{(x-t)^{1+\alpha} (y-\tau)^{1+\beta}} \cdot ii$$

Proof. Representation (8) itself is easily obtained by means of (2). Since $f \in H^{\lambda, \gamma}(Q)$, inequalities (9) are obvious. Estimate (10) is obtained by means of (5), i.e.

$$\psi(x, y) \leq C \left[x^{\lambda\theta} y^{(1-\theta)\gamma} + \alpha y^{(1-\theta)\gamma} \int_0^x \frac{dt}{(x-t)^{1-\theta\lambda}} + \beta x^{\theta\lambda} \int_0^y \frac{d\tau}{(y-\tau)^{1-(1-\theta)\gamma}} + \beta \alpha \int_0^x \int_0^y \frac{(x-t)^{\theta\lambda-1} dtd\tau}{(y-\tau)^{1-(1-\theta)\gamma}} \right]$$

It is easy to receive

$$\psi(x, y) \leq C x^{\theta\lambda} y^{(1-\theta)\gamma} \left[1 + \int_0^1 \frac{ds}{s^{1-\theta\lambda}} + \int_0^1 \frac{d\xi}{\xi^{1-(1-\theta)\gamma}} + \int_0^1 \int_0^1 \frac{s^{\theta\lambda-1} ds d\xi}{\xi^{1-(1-\theta)\gamma}} \right] \leq C_3 x^{\theta\lambda} y^{(1-\theta)\gamma}$$

Theorem 2. Let $f(x, y) \in H_0^{\lambda, \gamma}(Q)$, $\alpha < \lambda \leq 1$, $\beta < \gamma \leq 1$. Then the operator $D_{0+, 0+}^{\alpha, \beta}$ continuously maps $H_0^{\lambda, \gamma}(Q)$ into $H_0^{\lambda-\alpha, \gamma-\beta}(Q)$.

Proof. Since $f(x, y) \in H_0^{\lambda, \gamma}(Q)$, by (8) we have

$$\phi(x, y) = (D_{0+, 0+}^{\alpha, \beta} f)(x, y) = \psi(x, y)$$

Let $h > 0$; $x, x+h \in [0, b]$. We consider the difference

$$\begin{aligned} \psi(x+h, y) - \psi(x, y) &= \sum_{i=1}^{10} \Phi_k := \frac{\left(\Delta_{h, y}^{1,1} f \right) (0, 0)}{y^\beta (x+h)^\alpha} + \frac{\left(\Delta_{x, y}^{1,1} f \right) (0, 0)}{y^\beta} \left[\frac{1}{(x+h)^\alpha} - \frac{1}{x^\alpha} \right] + \\ &+ \frac{\alpha}{y^\beta} \int_0^x \frac{\left(\Delta_{h, y}^{1,1} f \right) (x, 0)}{(x+h-t)^{1+\alpha}} dt + \frac{\alpha}{y^\beta} \int_x^{x+h} \frac{\left(\Delta_{x+h-t, y}^{1,1} f \right) (t, 0)}{(x+h-t)^{1+\alpha}} dt + \frac{\beta}{(x+h)^\alpha} \int_0^y \frac{\left(\Delta_{h, y-\tau}^{1,1} f \right) (0, \tau)}{(y-\tau)^{1+\beta}} d\tau + \\ &+ \frac{\alpha}{y^\beta} \int_0^x \left(\Delta_{x-t, y}^{1,1} f \right) (t, 0) \left[(x+h-t)^{-1-\alpha} - (x-t)^{-1-\alpha} \right] dt + \beta \left[(x+h)^{-\alpha} - x^{-\alpha} \right] \int_0^y \frac{\left(\Delta_{x, y-\tau}^{1,1} f \right) (0, \tau)}{(y-\tau)^{1+\beta}} d\tau + \\ &+ \alpha \beta \int_0^x \int_0^y \frac{\left(\Delta_{h, y-\tau}^{1,1} f \right) (x, \tau) dtd\tau}{(x+h-t)^{1+\alpha} (y-\tau)^{1+\beta}} + \alpha \beta \int_x^{x+h} \int_0^y \frac{\left(\Delta_{x+h-t, y-\tau}^{1,1} f \right) (x, \tau) dtd\tau}{(x+h-t)^{1+\alpha} (y-\tau)^{1+\beta}} + \end{aligned}$$

$$+\alpha\beta \int_0^x \int_0^y \frac{\left(\Delta_{x-t, y-\tau}^{1,1} f\right)(x, \tau)}{(y-\tau)^{1+\beta}} \left[(x+h-t)^{-1-\alpha} - (x-t)^{-1-\alpha}\right] dt d\tau$$

(11)

Since $f(x, y) \in H_0^{\lambda, \gamma}(Q)$ we have

$$\begin{aligned} |\psi(x+h, y) - \psi(x, y)| &\leq \sum_{k=1}^{10} |\Phi_k| = C \left[\frac{h^\lambda}{y^\beta (x+h)^\alpha} + \frac{x^\lambda}{y^\beta} \left[\frac{1}{(x+h)^\alpha} - \frac{1}{x^\alpha} \right] + \frac{\alpha h^\lambda}{y^\beta} \int_0^x \frac{dt}{(x+h-t)^{1+\alpha}} + \right. \\ &+ \frac{\alpha}{y^\beta} \int_x^{x+h} (x+h-t)^{\lambda-1-\alpha} dt + \frac{h^{\theta\lambda} \beta}{(x+h)^\alpha} \int_0^y (y-\tau)^{(1-\theta)\gamma-1-\beta} d\tau + \frac{\alpha}{y^\beta} \int_0^x (x-t)^\lambda \left[\frac{1}{(x+h-t)^{1+\alpha}} - \frac{1}{(x-t)^{1+\alpha}} \right] dt + \\ &+ \beta x^{\theta\lambda} \left[(x+h)^{-\alpha} - x^{-\alpha} \right] \int_0^y \frac{d\tau}{(y-\tau)^{1+\beta-(1-\theta)\gamma}} + \alpha\beta h^{\theta\lambda} \int_0^x \int_0^y \frac{(y-\tau)^{(1-\theta)\gamma-1-\beta} dt d\tau}{(x+h-t)^{1+\alpha}} + \\ &+ \alpha\beta \int_x^{x+h} \int_0^y \frac{(y-\tau)^{(1-\theta)\gamma-1-\beta} dt d\tau}{(x+h-t)^{1+\alpha-\theta\lambda}} + \alpha\beta \int_0^x \int_0^y \frac{(x-t)^{\theta\lambda}}{(y-\tau)^{1+\beta-(1-\theta)\gamma}} \left[(x+h-t)^{-1-\alpha} - (x-t)^{-1-\alpha} \right] dt d\tau \end{aligned}$$

where

$$\int_0^y (y-\tau)^{(1-\theta)\gamma-1-\beta} d\tau < \infty$$

Using estimations G_1, G_2 of the proof of Lemma 1 and estimations I_1, I_2, I_3 of the proof of the Theorem 1, it is easily possible to receive an estimation

$$|\psi(x+h, y) - \psi(x, y)| \leq Ch^{\lambda-\alpha}$$

Rearranging symmetrically representation (11), we can similarly obtain that

$$|\psi(x, y+\eta) - \psi(x, y)| \leq C\eta^{\gamma-\beta}$$

References:

1. S.G. Samko, A.A. Kilbas and O.I. Marichev. Fractional Integrals and Derivatives. Theory and Applications. Gordon and Breach. Sci. Publ., N. York - London, 1993, 1012 pp. (Russian Ed. - Fractional Interals and Derivatives and Some of Their Applications, Nauka i Tekhnika, Minsk, 1987).

2. T.Mamatov Mixed fractional integration operators in Hölder spaces. «Science and World ». Volgograd, № 1 (1). 2013, P. 30-38
3. T.Mamatov, S. Samko. Mixed Fractional Integration Operators in Mixed Weighted Hölder Spaces. FC&AA. Vol.13, Num 3. 2010, p. 245-259
4. T. Mamatov. Weighted Zygmund estimates for mixed fractional integration. Case Studies Journal. Volume 7, Issue 5, 2018. p. 82-88
5. T. Mamatov. Mixed Fractional Integration In Mixed Weighted Generalized Hölder Spaces. Case Studies Journal. Volume 7, Issue 6. 2018, P. 61-68
6. T. Mamatov. Mixed Fractional Integration Operators in Hölder spaces. Monograph. LAPLAMBERT Academic Publishing. P. 73
7. T. Mamatov. Mixed Fractional Integration Operators in Mixed Weighted Hölder Spaces. Monograph. LAPLAMBERT Academic Publishing. P. 73